

Data based on first isolate per patient.

Daptomycin only routinely tested on Staphylococcus isolates recovered from blood or sterile body sites.

Nitrofurantoin only indicated for uncomplicated cystitis.

Streptococcus agalactiae (Group B Streptococci), Streptococcus pyogenes (Group A Streptococci), Streptococcus dysgalactiae, and Viridans group Streptococci only routinely tested if recovered from blood or sterile body sites.

Hospitals Included: Alton Memorial Hospital, Barnes-Jewish Hospital, Christian Hospital, Memorial Hospital, Parkland Health Center, St. Louis Children's Hospital

Ages: Adult (≥18 years)

Collection Setting: Inpatient, Outpatient, and Emergency Department

Antibiotic Class	Resistance Level	Antibiotic
85-100% of isolates susceptible		Trimethoprim-Sulfamethoxazole
		Tobramycin
		Piperacillin/Tazobactam
		Nitrofurantoin
60-84% of isolates susceptible		Minocycline
		Meropenem
0-59% of isolates susceptible		Levofloxacin
		Imipenem
Not routinely tested or insufficient data		Gentamicin
		Ciprofloxacin
		Ceftriaxone
		Cefazolidime
Intrinsically Resistant	#	Cefiderocol
		Cefepime
Inducibly Resistant	&	Cefazolin
		Aztreonam
		Ampicillin/Sulbactam
		Ampicillin

Nitrofurantoin only indicated for uncomplicated cystitis.

Ages: Adult (≥ 18 years)